

# Screening Checklist for Contraindications to Vaccines for Adults

Patient Name \_\_\_\_\_

Date of Birth \_\_\_\_ / \_\_\_\_ / \_\_\_\_  
month    day        year

**For patients:** The following questions will help us determine which vaccines you may be given today. If you answer “yes” to any question, it does not necessarily mean you should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

	Yes	No	Don't Know
1. Are you sick today?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Do you have allergies to medications, food, a vaccine component, or latex?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Have you ever had a serious reaction after receiving a vaccination?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Do you have a long-term health problem with heart disease, lung disease, asthma, kidney disease, metabolic disease (e.g., diabetes), anemia, or other blood disorder?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Do you have cancer, leukemia, HIV/AIDS, or any other immune system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. In the past 3 months, have you taken medications that affect your immune system, such as prednisone, other steroids, or anticancer drugs; drugs for the treatment of rheumatoid arthritis, Crohn's disease, or psoriasis; or have you had radiation treatments?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. Have you had a seizure or a brain or other nervous system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. During the past year, have you received a transfusion of blood or blood products, or been given immune (gamma) globulin or an antiviral drug?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. For women: Are you pregnant or is there a chance you could become pregnant during the next month?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. Have you received any vaccinations in the past 4 weeks?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Form completed by \_\_\_\_\_ Date \_\_\_\_\_

Form reviewed by \_\_\_\_\_ Date \_\_\_\_\_

Did you bring your immunization record card with you? Yes  No

# Hepatitis A Vaccine

## What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See [www.imzmniza.org/via](http://www.imzmniza.org/via)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.imzmniza.org/via](http://www.imzmniza.org/via)

### 1 What is hepatitis A?

Hepatitis A is a serious liver disease caused by the hepatitis A virus (HAV). HAV is found in the stool of people with hepatitis A.

It is usually spread by close personal contact and sometimes by eating food or drinking water containing HAV. A person who has hepatitis A can easily pass the disease to others within the same household.

Hepatitis A can cause:

- “flu-like” illness
- jaundice (yellow skin or eyes, dark urine)
- severe stomach pains and diarrhea (children)

People with hepatitis A often have to be hospitalized (up to about 1 person in 5).

Adults with hepatitis A are often too ill to work for up to a month.

Sometimes, people die as a result of hepatitis A (about 3–6 deaths per 1,000 cases).

Hepatitis A vaccine can prevent hepatitis A.

### 2 Who should get hepatitis A vaccine and when?

#### WHO

*Some people should be routinely vaccinated with hepatitis A vaccine:*

- All children between their first and second birthdays (12 through 23 months of age).
- Anyone 1 year of age and older traveling to or working in countries with high or intermediate prevalence of hepatitis A, such as those located in Central or South America, Mexico, Asia (except Japan), Africa, and eastern Europe. For more information see [www.cdc.gov/travel](http://www.cdc.gov/travel).
- Children and adolescents 2 through 18 years of age who live in states or communities where routine vaccination has been implemented because of high disease incidence.
- Men who have sex with men.
- People who use street drugs.
- People with chronic liver disease.

- People who are treated with clotting factor concentrates.
- People who work with HAV-infected primates or who work with HAV in research laboratories.
- Members of households planning to adopt a child, or care for a newly arriving adopted child, from a country where hepatitis A is common.

*Other people might get hepatitis A vaccine in certain situations (ask your doctor for more details):*

- Unvaccinated children or adolescents in communities where outbreaks of hepatitis A are occurring.
- Unvaccinated people who have been exposed to hepatitis A virus.
- Anyone 1 year of age or older who wants protection from hepatitis A.

Hepatitis A vaccine is not licensed for children younger than 1 year of age.

#### WHEN

For children, the first dose should be given at 12 through 23 months of age. Children who are not vaccinated by 2 years of age can be vaccinated at later visits.

For others at risk, the hepatitis A vaccine series may be started whenever a person wishes to be protected or is at risk of infection.

For travelers, it is best to start the vaccine series at least one month before traveling. (Some protection may still result if the vaccine is given on or closer to the travel date.)

Some people who cannot get the vaccine before traveling, or for whom the vaccine might not be effective, can get a shot called immune globulin (IG). IG gives immediate, temporary protection.

Two doses of the vaccine are needed for lasting protection. These doses should be given at least 6 months apart.

Hepatitis A vaccine may be given at the same time as other vaccines.



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

**3****Some people should not get hepatitis A vaccine or should wait.**

- Anyone who has ever had a severe (life threatening) allergic reaction to a previous dose of hepatitis A vaccine should not get another dose.
- Anyone who has a severe (life threatening) allergy to any vaccine component should not get the vaccine.
- **Tell your doctor if you have any severe allergies, including a severe allergy to latex.** All hepatitis A vaccines contain alum, and some hepatitis A vaccines contain 2-phenoxyethanol.
- Anyone who is moderately or severely ill at the time the shot is scheduled should probably wait until they recover. Ask your doctor. People with a mild illness can usually get the vaccine.
- Tell your doctor if you are pregnant. Because hepatitis A vaccine is inactivated (killed), the risk to a pregnant woman or her unborn baby is believed to be very low. But your doctor can weigh any theoretical risk from the vaccine against the need for protection.

**4****What are the risks from hepatitis A vaccine?**

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of hepatitis A vaccine causing serious harm, or death, is extremely small.

Getting hepatitis A vaccine is much safer than getting the disease.

**Mild problems**

- soreness where the shot was given (*about 1 out of 2 adults, and up to 1 out of 6 children*)
- headache (*about 1 out of 6 adults and 1 out of 25 children*)
- loss of appetite (*about 1 out of 12 children*)
- tiredness (*about 1 out of 14 adults*)

If these problems occur, they usually last 1 or 2 days.

**Severe problems**

- serious allergic reaction, within a few minutes to a few hours after the shot (*very rare*).

**5****What if there is a serious reaction?****What should I look for?**

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

**What should I do?**

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at [www.vaers.hhs.gov](http://www.vaers.hhs.gov), or by calling 1-800-822-7967.

*VAERS is only for reporting reactions. They do not give medical advice.*

**6****The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation).

**7****How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call 1-800-232-4636 (1-800-CDC-INFO) or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines)

Vaccine Information Statement (Interim)

**Hepatitis A Vaccine**

10/25/2011

42 U.S.C. § 300aa-26

